

Memo

Use of the paediatric Pfizer COVID-19 vaccine in 5-11 year-olds – second dose and dosing interval: COVID-19 Vaccine Technical Advisory Group (CV TAG) recommendations

Date: 16 February 2022

To: Dr Ashley Bloomfield, Director-General of Health

Copy to: Astrid Koornneef, Director, National Immunisation Programme
Allison Bennett, Manager, System Enablers, System Strategy and Policy
Dr Caroline McElnay, Director of Public Health

From: Dr Ian Town, Chief Science Advisor

For your: Consideration

Purpose of report

1. To outline the COVID-19 Vaccine Technical Advisory Group's (CV TAG) advice about the administration of a second dose of the paediatric Pfizer vaccine and the interval between the first and second doses of the COVID-19 vaccine for 5–11 year-olds.
2. This report also provides an update on international and local safety data.

Background and context

3. Vaccination of 5–11-year-olds in New Zealand is now underway. The approved COVID-19 paediatric Pfizer vaccine being used has a lower dose (10 µg) and a smaller volume (0.2 mL) than the adult vaccine and is administered using a smaller needle. As at 13 February 2022, 214,857 (45%) of 5–11-year-olds had received their first dose in New Zealand. [1] Only 26% of Māori 5-11 year-olds and 36% of Pacific 5-11 year-olds have received their first dose. To be fully immunised against COVID-19, a child needs to receive two doses of the paediatric vaccine.
4. In December 2021, CV TAG recommended that two doses of the paediatric Pfizer vaccine be offered to all 5-11 year-olds in Aotearoa New Zealand, with an 8-week interval between doses (Appendix 1, *Decision to Use the Pfizer mRNA COVID-19 vaccine for children aged 5-11 years: COVID-19 Vaccine Technical Advisory Group (CV TAG) recommendations*). It was also indicated that in February 2022, CV TAG would assess the latest data and provide updated recommendations prior to any second doses being given to this age group in New Zealand.

Safety data for the Pfizer vaccine in 5–11-year-olds

5. A randomised clinical trial to assess the safety, immunogenicity, and efficacy of the Pfizer vaccine in 5-11 year-olds of two doses administered three weeks apart reported more local reactions and systemic events than placebo recipients. [2] The reactions and events reported were generally mild to moderate, lasting 1 to 2 days. Injection-site pain was the most common local reaction, occurring in 71 to 74% of Pfizer recipients. Severe injection-site pain after the first or second dose was reported in 0.6% of Pfizer recipients and in no placebo recipients.
6. In the clinical trial, fatigue and headache were the most frequently reported systemic events (0.9%), headache (0.3%), chills (0.1%), and muscle pain (0.1%) were also reported after the first or second dose of Pfizer. [2] Frequencies of fatigue, headache, and chills were similar among Pfizer and placebo recipients after the first dose and were more frequent among Pfizer recipients than among placebo recipients after the second dose.
7. No vaccine-related serious adverse events were noted in the clinical trial; however, the trial was too small to detect rare side effects such as myocarditis. [2] Three serious adverse events were reported from two participants (postinjury abdominal pain and pancreatitis in a placebo recipient and arm fracture in a Pfizer recipient), however, none of these were related to the vaccine or placebo. No deaths or adverse events leading to withdrawal were reported. There were no cases of severe COVID-19 or Multisystem Inflammatory Syndrome in Children (MIS-C)—a condition associated with COVID-19 where body parts can become inflamed. Lymphadenopathy was reported in ten Pfizer recipients (0.9%) and one placebo recipient (0.1%). No myocarditis, pericarditis, hypersensitivity, or anaphylaxis in Pfizer recipients was reported. Rashes in four Pfizer recipients (observed on the arm, torso, face, or body, with no consistent pattern) were related to vaccination; the rashes were mild and self-limiting, and onset was typically 7 days or more after vaccination.
8. Real-world safety data has been collected from over 8 million doses of the Pfizer vaccine administered to children aged 5–11 years in the United States. These data have been collected in the Vaccine Adverse Event Reporting System (VAERS), a national passive vaccine safety surveillance system, and through V-safe, a voluntary smartphone-based safety surveillance system for adverse events after COVID-19 vaccination. [3] From November 3 to December 19, 2021, VAERS received and processed 4,249 reports of adverse events for children aged 5–11 years who received Pfizer COVID-19 vaccine. Overall, among VAERS reports for children aged 5–11 years who received the Pfizer vaccine, approximately 97% were non-serious. The most commonly reported conditions among the 100 reports of serious events were fever (29.0%), vomiting (21.0%), and increased troponin-(15.0%). Among 12 serious reports of seizure, five children experienced new-onset seizures. Among 15 preliminary reports of myocarditis identified during the analytic period, 11 met the case definition for myocarditis. VAERS received two reports of death both of whom had complicated medical histories and were in fragile health before vaccination. None of the data suggested a causal association between death and vaccination. In V-safe, fever was found to be more frequently reported in 5-11 year-olds after dose 2 (4,001: 13.4%) than dose 1 (3,350; 7.9%) among 42,504 recipients of dose 1 and 29,899 recipients of dose 2. Overall, systemic reactions after dose 2 among registrants aged 5-11 years were less frequent than among children aged 12-15 years. Fourteen registrants aged 5–11 years received hospital care after vaccination. Information regarding reason for hospitalisation was available for five children

and included appendicitis (two), vomiting and dehydration (one), respiratory infection (one), and retropharyngeal cellulitis (one).

Reporting of Adverse Events following Vaccination in New Zealand

9. In New Zealand, preliminary unpublished data from Medsafe indicates that there have been 352 adverse events following immunisation (AEFIs) reported from 17 January to 30 January 2022 in children aged 5-11 who received the approved COVID-19 paediatric Pfizer vaccine. Of these, 96.9% (341) reports were classified as non-serious. A small number of individuals (10) reported that an AEFI required emergency care and one AEFI case was admitted to hospital for observation (no evidence of myocarditis despite reporting chest discomfort). Of these 11 cases, six were reported as recovered or recovering, one was ongoing, and four had an unknown outcome. Chest discomfort was the most frequently reported reaction (6), followed by vasovagal reaction (4), and there was one case of anaphylaxis (Brighton criteria level 4).
10. Medsafe is in regular contact with other regulators and have noted that to date nothing of concern has been drawn to their attention regarding the safety profile of the paediatric Pfizer vaccine.

Rationale for an 8-week interval

11. The manufacturer's recommended schedule for the paediatric Pfizer vaccine is 2 doses, 3 weeks apart.
12. Research conducted in adults into extending the dosing interval (e.g., to 8 weeks or longer) has shown that longer intervals between the first and second Pfizer dose can lead to higher humoral and cellular immune responses, improved vaccine effectiveness, and potentially a longer duration of protection compared with the standard interval. [4-7] In addition, data from adults show that an extended dosing interval may also reduce the risk of myocarditis and pericarditis after vaccination. [8]
13. Extended dosing intervals has not yet been studied in children, but it is expected that similar effects would be observed to those after extended dosing intervals in adults, such as improved immunogenicity and the potential for a lower risk of serious side effects. The recommendation for an 8-week interval between doses is consistent with other international advisory groups, such as in the UK, Canada, and Australia. [9-11] In addition, a longer interval between doses would allow more time to continue monitoring international safety data as it emerges.

Priority groups for children aged 5-11 years

14. Māori and Pacific children have been disproportionately affected in this pandemic. For community-acquired cases up to 11 February 2022, Māori made up 45.7% of total cases in 5- to 11-year-olds, and Pacific children have made up 28.7% of cases among 5- to 11-year-olds. Of these cases, a total of ten have been hospitalised, with Māori and Pacific children combined making up 90% of these cases. As noted above, in the vaccine rollout for 5-11-year-olds, fewer Māori and Pacific children have been vaccinated than other ethnicities. Prioritisation of Māori and Pacific children remains important, and the emphasis should be to get the first dose administered to as many as children as possible.

15. Children living with pre-existing conditions or comorbidities, from disadvantaged backgrounds, or those living within a lower socioeconomic status have a greater risk of severe disease from COVID-19. [12-16]
16. Starship Child Health has listed risk factors for COVID-19 disease [17] that may be used as guidance for prioritising children with high-risk pre-existing conditions. The current list of risk factors includes children with:
 - Chronic lung disease including bronchiectasis, cystic fibrosis, BiPAP for OSA
 - Non-repaired congenital heart disease, acquired heart disease or congestive heart failure
 - Poorly controlled asthma (regular symptoms occurring in a usual week that affect the patient's quality of life and includes anyone with an admission in the last 2 years or anyone with 2 or more courses of steroids in the last two years)
 - Obesity (BMI \geq 95th centile for age)
 - Diabetes (insulin-dependent)
 - Chronic kidney disease (GFR $<$ 15 ml/min/1.73m²)
 - Severe cerebral palsy (or neurodevelopmental disorder)
 - Complex genetic, metabolic disease or multiple congenital anomalies.
17. Children in other recognised clinical risk groups who are at higher risk of severe COVID-19 should also include those who are a household contact of someone who is immunosuppressed (defined as those who expect to share living accommodation on most days (and therefore for whom continuing close contact is unavoidable) with individuals of any age who are immunosuppressed).

Recommendations

18. CV TAG met on 1, 8, and 15 February 2022 to consider guidance on administering a second dose of the vaccine and the interval between doses for 5–11-year-olds.
19. CV TAG noted:
 - a. **The direct and indirect impacts on children.** Children who have COVID-19 will commonly have few or only mild respiratory symptoms. COVID-19 in this age group is rarely severe or fatal, [18, 19] and the rate of severe COVID-19 disease in this age group is the lowest of any age group. However, there is a very small but real risk of MIS-C (described above) at this age which has occurred more frequently among ethnic minorities in the US. [12, 20] A very small proportion of children also experience persistent illness and ongoing symptoms, though evidence about its incidence is limited.
 - b. Children living with pre-existing conditions or comorbidities, from disadvantaged backgrounds, or those living within a lower socioeconomic status have a greater risk of severe disease from COVID-19. [12-15]
 - c. Even though the direct effects of infection are generally less severe in children, this should not diminish the significance for those who have experienced worse outcomes. [19] Alongside the direct risks and impacts to health and individuals, COVID-19 also has indirect impacts for children on mental health, wellbeing, education and social development, and these are worsened by lockdowns and school closures. [12, 21-23]
 - d. **Children do play a role in transmission however it is significantly smaller than for adults.** Transmission within education settings has occurred but is limited and is more

likely to occur between adults. [24-26] Transmission in households is much more common. [27, 28] The benefit of vaccination on onward transmission in households could be lower than in other settings due to the ongoing and close nature of exposure. [29, 30] but this is not confirmed. The effect of vaccination of children on household transmission is unknown.

- e. **There are a number of equity considerations which are important to consider:**
- i. Māori and Pacific children have been disproportionately affected in the current outbreak. To 11 February 2022, Māori made up 45.7% of cases in 5-11 year-olds, and Pacific children have made up 28.7% of cases among 5-11 year-olds.
 - ii. Māori and Pacific adults are at greater risk of COVID-19 hospitalisation and severe disease. Māori and Pacific adults have respectively 2.5-fold and 3-fold higher odds of being hospitalised compared to non-Māori, and Māori are likely to spend 4.9 days longer in hospital. [31, 32]
 - iii. Māori and Pacific children are more likely to live in multigenerational families housed in overcrowded conditions, increasing the risk of transmission. The younger age structure of the Māori population also means that a larger proportion are currently unable to be vaccinated and remain susceptible to infection and transmission, with a risk of onwards transmission to whānau and communities, [33, 34] though the risk of transmission from children is lower than from adults.
- f. **The paediatric formulation of the vaccine has been approved for emergency use and rolled out in the USA, Canada, and Israel.** The Advisory Committee on Immunization Practices (ACIP) made an interim recommendation for the emergency use of the Pfizer vaccine in children aged 5-11 years in the United States for prevention of COVID-19. [19, 35] This was unanimously supported by the Committee. In making this recommendation, ACIP considered the importance of COVID-19 as a public health problem, as well as benefits and harms, parents' values and preferences, acceptability, feasibility, resource use, and equity for use of the vaccine among children. [19, 35] ACIP additionally stated: "children from racial and ethnic minority groups have experienced a disproportionately high incidence of COVID-19 as well as secondary impacts of the COVID-19 pandemic such as reduced in-person learning". [19] These comments have high relevance for New Zealand given the similar effects of the pandemic on Māori and Pacific Peoples as described above.
- g. **In Australia, the Therapeutic Goods Administration (TGA)** provisionally approved the Pfizer vaccine as safe and effective for use among this age group on 5 December 2021. [36] ATAGI recommends all 5–11-year-olds be vaccinated with an 8-week interval between doses, and that those at risk of severe disease, Aboriginal and Torres Strait Islanders, and children in crowded conditions or outbreak areas be prioritised. [37]
- h. **On dosing intervals,** there are no data available about extending the interval between doses of the paediatric formulation of the Pfizer vaccine, however, emerging data in adults suggests that the immune response is likely improved by extending the dosing interval. [4-7] This is consistent with basic principles of vaccinology and immunology which suggests that immune responses are generally better with longer intervals. There may also be a connection between shorter intervals and increased reactogenicity or adverse events, and one pre-print paper on individuals aged 12 and over has shown

a statistically significant increase in myocarditis if the second dose was given at a shorter interval of less than 30 days. [8] Australia, Canada, and the UK have recommended an 8-week interval between doses for 5-11 year-olds, noting this may improve immunogenicity and reduce side effects. Having a longer interval would also allow more time to monitor international safety data.

- i. **On vaccine requirements**, there is a significant risk that use of vaccination mandates or certificates in this age group will result in exclusion and an inability to fully participate in schooling and extracurricular activities. This is likely to inequitably impact communities who are already experiencing disadvantage and where current vaccine coverage is poor. Concerns regarding possible stigmatisation and exclusions could be addressed in ways that do not necessarily influence the decision to use. For example, there could be a policy decision that children cannot be denied access to locations/events on the basis of vaccination status, which could be operationalised by not issuing vaccine certificates for this age group.
- j. **On safety of the paediatric vaccine**, real-world data on the rollout of the vaccine to 5–11-year-olds have reported nothing of concern to date.

20. **CV TAG recommended that:**

- a. A second dose of the paediatric Pfizer vaccine be offered to all 5–11 year-olds in Aotearoa New Zealand, with a minimum 8-week interval between doses.
- b. Māori and Pacific children, children with high-risk pre-existing conditions, and children living with vulnerable people should continue to be prioritised for vaccination.

21. CV TAG will continue to monitor all relevant information (including safety data) and will update their recommendations as information becomes available.

Ian G Town

Dr Ian Town
**Chief Science Advisor and
Chair of the COVID-19 Vaccine Technical Advisory Group**

References

1. New Zealand Ministry of Health. *COVID-19: Vaccine data*. 13 February 2022; Available from: <https://www.health.govt.nz/covid-19-novel-coronavirus/covid-19-data-and-statistics/covid-19-vaccine-data>.
2. Walter, E.B., et al., *Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age*. *New England Journal of Medicine*, 2021.
3. Hause, A.M., et al., *COVID-19 Vaccine Safety in Children Aged 5–11 Years — United States, November 3–December 19, 2021*. *MMWR. Morbidity and Mortality Weekly Report*, 2021. **70**(5152): p. 1755-1760.
4. Hall, V.G., et al., *Delayed-interval BNT162b2 mRNA COVID-19 vaccination enhances humoral immunity and induces robust T cell responses*. *Nature Immunology*, 2022.
5. Parry, H., et al., *Extended interval BNT162b2 vaccination enhances peak antibody generation*. *npj Vaccines*, 2022. **7**(1): p. 14.
6. Tauzin, A., et al., *Strong humoral immune responses against SARS-CoV-2 Spike after BNT162b2 mRNA vaccination with a 16-week interval between doses*. *Cell Host & Microbe*, 2022. **30**(1): p. 97-109.e5.
7. Payne, R.P., et al., *Immunogenicity of standard and extended dosing intervals of BNT162b2 mRNA vaccine*. *Cell*, 2021. **184**(23): p. 5699-5714.e11.
8. Buchan, S.A., et al., *Epidemiology of myocarditis and pericarditis following mRNA vaccines in Ontario, Canada: by vaccine product, schedule and interval*. *medRxiv*, 2021: p. 2021.12.02.21267156.
9. UK Government. *JCVI statement on COVID-19 vaccination of children and young people: 22 December 2021*. 22 December 2021; Available from: <https://www.gov.uk/government/publications/jcvi-update-on-advice-for-covid-19-vaccination-of-children-and-young-people/jcvi-statement-on-covid-19-vaccination-of-children-and-young-people-22-december-2021>.
10. Government of Canada. *NACI updated recommendations on the use of COVID-19 vaccines in children 5 to 11 years of age*. January 25, 2022; Available from: <https://www.canada.ca/en/public-health/services/immunization/national-advisory-committee-on-immunization-naci/updated-recommendations-use-covid-19-vaccines-children-5-11-years-age.html>.
11. Australian Government Department of Health. *ATAGI recommendations on Pfizer COVID-19 vaccine use in children aged 5 to 11 years*. 10 December 2021; Available from: <https://www.health.gov.au/resources/publications/ataqi-recommendations-on-pfizer-covid-19-vaccine-use-in-children-aged-5-to-11-years>.
12. Murdoch Children's Research Institute (MCRI). *Research Brief: COVID-19 and Child and Adolescent Health*. Research Brief 2021 13 September 2021 [cited 2021 16 November]; Version 1:[Available from: <https://www.mcri.edu.au/sites/default/files/media/documents/covid-19-and-child-and-adolescent-health-140921.pdf>].
13. Tsankov, B.K., et al., *Severe COVID-19 Infection and Pediatric Comorbidities: A Systematic Review and Meta-Analysis*. *Int J Infect Dis*, 2021. **103**: p. 246-256.
14. Ng, W.H., et al., *Comorbidities in SARS-CoV-2 Patients: a Systematic Review and Meta-Analysis*. *mBio*, 2021. **12**(1).
15. Mukkada, S., et al., *Global characteristics and outcomes of SARS-CoV-2 infection in children and adolescents with cancer (GRCCC): a cohort study*. *Lancet Oncol*, 2021. **22**(10): p. 1416-1426.
16. Shi, Q., et al., *Risk factors for poor prognosis in children and adolescents with COVID-19: A systematic review and meta-analysis*. *EClinicalMedicine*, 2021. **41**.
17. Starship Child Health. *COVID-19 disease in children*. 24 January 2022; Available from: <https://starship.org.nz/guidelines/covid-19-disease-in-children/>.

18. Ludvigsson, J.F., *Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults*. *Acta Paediatr*, 2020. **109**(6): p. 1088-1095.
19. Woodworth, K.R., et al., *The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-BioNTech COVID-19 Vaccine in Children Aged 5–11 Years — United States, November 2021*. *MMWR Morb Mortal Wkly Rep*, 2021. **70**: p. 1579–1583.
20. Centers for Disease Control and Prevention (CDC). *For Parents: Multisystem Inflammatory Syndrome in Children (MIS-C) associated with COVID-19*. Multisystem Inflammatory Syndrome 2021 20 September 2021 [cited 2021 16 November]; Available from: <https://www.cdc.gov/mis/mis-c.html>.
21. Health, A.I.o. and Welfare, *COVID-19 and the impact on young people*. 2021, AIHW: Canberra.
22. Brown, N., Te Riele, K., Shelley, B. & Woodroffe, J., *Learning at home during COVID-19: Effects on vulnerable young Australians. Independent Rapid Response Report.*, P.U.C.f.E. Attainment., Editor. 2020, University of Tasmania: Hobart.
23. Roubinov, D., N.R. Bush, and W.T. Boyce, *How a Pandemic Could Advance the Science of Early Adversity*. *JAMA Pediatrics*, 2020. **174**(12): p. 1131-1132.
24. European Centre for Disease Prevention and Control. *Questions and answers on COVID-19: Children aged 1 – 18 years and the role of school settings*. 2021 8 September 2021; Available from: <https://www.ecdc.europa.eu/en/covid-19/questions-answers/questions-answers-school-transmission>.
25. Falk A, B.A., Falk P, Steffen S, Wallace Z, Høeg TB. , *COVID-19 Cases and Transmission in 17 K–12 Schools — Wood County, Wisconsin, August 31–November 29, 2020*. *MMWR Morb Mortal Wkly Rep*, 2021. **70**: p. 136–140.
26. Ismail, S.A., et al., *SARS-CoV-2 infection and transmission in educational settings: a prospective, cross-sectional analysis of infection clusters and outbreaks in England*. *The Lancet Infectious Diseases*, 2021. **21**(3): p. 344-353.
27. Dattner, I., et al., *The role of children in the spread of COVID-19: Using household data from Bnei Brak, Israel, to estimate the relative susceptibility and infectivity of children*. *PLOS Computational Biology*, 2021. **17**(2): p. e1008559.
28. Li, W., et al., *Characteristics of Household Transmission of COVID-19*. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*, 2020. **71**(8): p. 1943-1946.
29. Shah, A.S.V., et al. *Effect of vaccination on transmission of COVID-19: an observational study in healthcare workers and their households*. 21 March 2021; Available from: <https://doi.org/10.1101/2021.03.11.21253275>.
30. Harris, R.J., et al., *Effect of Vaccination on Household Transmission of SARS-CoV-2 in England*. *New England Journal of Medicine*, 2021.
31. Steyn, N., Binny, R. N., Hannah, K., Hendy, S. C., James, A., Lustig, A., Ridings, K., Plank, M. J., Sporle, A., *Māori and Pacific people in New Zealand have a higher risk of hospitalisation for COVID-19*. *New Zealand Medical Journal*, 2021. **134**(1538).
32. Steyn, N., et al., *Estimated inequities in COVID-19 infection fatality rates by ethnicity for Aotearoa New Zealand*. *New Zealand Medical Journal*, 2020. **133**(1521): p. 28-39.
33. McLeod, M., et al., *COVID-19: we must not forget about Indigenous health and equity*. *Australian and New Zealand Journal of Public Health*, 2020. **44**(4): p. 253-256.
34. Johnson, A., P. Howden-Chapman, and S. Eaquad, *A stocktake of New Zealand's housing: February 2018*. 2018: Ministry of Business, Innovation and Employment (New Zealand).
35. Advisory Committee on Immunization Practices (ACIP), *ACIP Evidence to Recommendations for Use of Pfizer-BioNTech COVID-19 Vaccine under an Emergency Use Authorization*. 2021.

36. Department of Health. *TGA provisionally approves Pfizer COVID-19 vaccine for 5 to 11-year-olds*. 2021 5 December 2021; Available from: <https://www.health.gov.au/ministers/the-hon-greg-hunt-mp/media/tga-provisionally-approves-pfizer-covid-19-vaccine-for-5-to-11-year-olds>.
37. ATAGI. *ATAGI expanded guidance on acute major medical conditions that warrant a temporary medical exemption relevant for COVID-19 vaccines*. 2021; Available from: <https://www.health.gov.au/sites/default/files/documents/2021/09/atagi-expanded-guidance-on-temporary-medical-exemptions-for-covid-19-vaccines.pdf>.
38. Polack, F.P., et al., *Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine*. *New England Journal of Medicine*, 2020. **383**(27): p. 2603-2615.
39. Frenck, R.W., et al., *Safety, Immunogenicity, and Efficacy of the BNT162b2 Covid-19 Vaccine in Adolescents*. *New England Journal of Medicine*, 2021.
40. NSW Health. *EPIDEMIOLOGICAL WEEK 45 ENDING 13 NOVEMBER 2021. COVID-19 WEEKLY SURVEILLANCE IN NSW 2021* [cited 2021 2 December]; Available from: <https://www.health.nsw.gov.au/Infectious/covid-19/Documents/covid-19-surveillance-report-20211113.pdf>.
41. RNZ. *Concern over Covid-19 vaccine for children: 'Protective instinct kicks in'*. 2021 9 December 2021; Available from: <https://www.rnz.co.nz/news/national/457592/concern-over-covid-19-vaccine-for-children-protective-instinct-kicks-in>.
42. Skowronski, D.M., et al., *Two-dose SARS-CoV-2 vaccine effectiveness with mixed schedules and extended dosing intervals: test-negative design studies from British Columbia and Quebec, Canada*. *medRxiv*, 2021: p. 2021.10.26.21265397.
43. Flaxman, A., et al., *Reactogenicity and immunogenicity after a late second dose or a third dose of ChAdOx1 nCoV-19 in the UK: a substudy of two randomised controlled trials (COV001 and COV002)*. *Lancet*, 2021. **398**(10304): p. 981-990.
44. Buchan, S., et al., *Epidemiology of myocarditis and pericarditis following mRNA vaccines in Ontario, Canada: by vaccine product, schedule and interval*. 2021, *medRxiv*.

**Appendix 1 – Decision to Use the Pfizer mRNA COVID-19 vaccine for children aged 5-11 years:
COVID-19 Vaccine Technical Advisory Group (CV TAG) recommendations**

Memo

Decision to use the Pfizer mRNA COVID-19 vaccine for children aged 5-11 years: COVID-19 Vaccine Technical Advisory Group (CV TAG) recommendations

Date: 15 December 2021

To: Dr Ashley Bloomfield, Director-General of Health

Copy to: Astrid Koornneef, Director of National Operations, COVID Vaccine Immunisation Programme
Allison Bennett, Manager, System Enablers, System Strategy and Policy
Dr Caroline McElnay, Director of Public Health

From: Dr Ian Town, Chief Science Advisor and Chair of CV TAG

For your: Information

Purpose of report

1. To summarise the CV TAG recommendations on the decision to use the paediatric formulation of the Pfizer mRNA COVID-19 vaccine ('the Pfizer vaccine') for children who are 5 to 11 years of age.

Background and context

2. In February 2021, CV TAG advice was sought for use of the Pfizer COVID-19 vaccine in people who were 16 years and over, following Medsafe approval. Cabinet agreed that the COVID-19 Immunisation Programme proceed with the roll out of the Pfizer vaccine, and this has been underway since February.
3. In August 2021, CV TAG confirmed support to extend the age of people who can receive the Pfizer vaccine to 12- to 15-year-olds, noting that this would likely lead to a reduction in school closures and disruption to education, and contribute to equitable vaccination coverage in Māori and Pacific peoples.
4. Medsafe is assessing an application submitted by Pfizer for the use of a paediatric formulation of the vaccine in 5- to 11-year-olds within New Zealand. The CV TAG recommendations presented here are subject to Medsafe approval and any listed clinical conditions.

5. The Ministry's Policy team has sought clinical and scientific advice from CV TAG on the use of the Pfizer vaccine for children who are 5- to 11-years of age. This advice will be considered as part of the Decision to Use Framework, and alongside policy considerations for the sequencing of the COVID-19 Immunisation Programme.

The COVID-19 vaccine in 5- to 11-year-olds

Phase 2/3 trial findings

6. One phase 2/3 randomised control trial was conducted to assess the safety, immunogenicity, and efficacy of two doses of the Pfizer vaccine administered 21 days apart in children aged 6 months to 11 years, with findings published for 5- to 11-year-olds to date [2].
7. In the phase 2/3 trial, participants were randomly assigned in a 2:1 ratio to receive two doses of either the Pfizer vaccine at 10 µg (a lower dose than the 30 µg used in older age groups), or a placebo. A total of 2268 children were assigned to receive the Pfizer vaccine (1517 children) or placebo (751 children) [2].
8. The trial was run across 81 sites in the US, Spain, Finland and Poland. Overall, 52% were male, 79% were White, 6% were Black, 6% were Asian, and 21% were Hispanic or Latinx. The mean age was 8.2 years; 20% of children had coexisting conditions (including 12% with obesity and approximately 8% with asthma), and 9% were SARS-CoV-2-positive at baseline. Demographic characteristics were similar between the 5- to 11-year-old and 16- to 25-year-old Pfizer recipients who were included in the immuno-bridging subset, apart from younger age and the percentage of Black and Hispanic or Latinx in the 5- to 11-year-old group (6% and 18%, respectively) being lower than in the 16- to 25-year-old group (12% and 36%, respectively) [2].

Safety and reactogenicity

9. In the 5- to 11-year-olds, as in other age groups, the Pfizer vaccine had a favourable safety profile, with side effects generally comparable to those observed in 16- to 25-year-olds who received the standard 30 µg doses [2].
10. Safety evaluations included assessment of reactogenicity events reported by a parent or guardian using an electronic diary for 7 days after each dose. Data on unsolicited adverse events, including confirmed diagnoses of myocarditis or pericarditis, were collected from the first dose through 1 month after the second dose. Data on serious adverse events will be collected from the first dose through 6 months after the second dose [2]. At data cut-off, the median follow-up was 2.3 months [2].
11. Most local reactions were mild to moderate, lasting 1-2 days. Injection-site pain was the most common local reaction, occurring in 71-74% of Pfizer recipients. Fatigue and headache were the most frequently reported systemic events. In general, systemic events were reported more frequently after the second dose than first dose. As compared with adults and adolescents in the pivotal trial, 5- to 11-year-olds reported a higher incidence of injection-site redness (15 to 19%, vs. 5 to 7%) and swelling (10 to 15%, vs. 5 to 8%), but a generally lower incidence of systemic events, including fever (3 to 7%, vs. 1 to 20%) and chills (5 to 10%, vs. 6 to 42%) [2, 38, 39].

12. From the first dose through to one month after the second dose, adverse events were reported by 10.9% of Pfizer recipients and 9.2% of placebo recipients. Slightly more Pfizer recipients (3.0%) than placebo recipients (2.1%) reported adverse events that were considered by the investigators to be related to the vaccine or placebo [2].
13. No vaccine-related serious adverse events were noted in the clinical trial; however, the trial was too small to detect rare side effects such as myocarditis. Three serious adverse events were reported from two participants (postinjury abdominal pain and pancreatitis in a placebo recipient and arm fracture in a Pfizer recipient). None of these were considered to be related to the vaccine or placebo. No deaths or adverse events leading to withdrawal were reported [2]. No cases of severe COVID-19 or Multisystem Inflammatory Syndrome in Children (MIS-C) were reported—a condition associated with COVID-19 where body parts can become inflamed [2, 20]. Lymphadenopathy was reported in 10 Pfizer recipients (0.9%) and 1 placebo recipient (0.1%). No myocarditis, pericarditis, hypersensitivity, or anaphylaxis in Pfizer recipients was reported. Four rashes in Pfizer recipients (observed on the arm, torso, face, or body, with no consistent pattern) were considered to be related to vaccination; the rashes were mild and self-limiting, and onset was typically 7 days or more after vaccination [2].
14. No safety data are yet available from the large-scale roll out of the Pfizer vaccine to 5- to 11-year-olds in the USA, though will likely be available by late December 2021 or early January 2022.

Immunogenicity and efficacy

15. Immune responses in the single clinical trial conducted were assessed one month after the second dose of the Pfizer vaccine were equivalent to those in 16- to 25-year-olds. Children aged 5-11 receiving two 10 µg doses had a similar, statistically non-inferior, neutralising antibody responses with a geometric mean titre (GMT) of 1,197.6 (95% CI: 1,106.1, 1,296.6) vs 1,146.5 (95% CI: 1,045.5, 1,257.2) for ages 16-25 [2].
16. Vaccine efficacy against symptomatic NAAT-confirmed COVID-19 at 7 days or more after the second dose (to a median follow up of 2.3 months at data cut-off) was assessed. Among participants without evidence of previous SARS-CoV-2 infection, symptomatic COVID-19 was reported in three recipients of the Pfizer vaccine and in 16 placebo recipients, producing a vaccine efficacy of 90.7% (95% CI, 67.7 to 98.3) [2].

CV TAG Recommendations

17. CV TAG discussed the use of the Pfizer COVID-19 vaccine in children aged 5-11 years at meetings between October and December 2021 and consulted with Māori paediatricians and Māori general practitioners at two meetings in December 2021.¹
18. CV TAG noted:

¹ CV TAG discussed use of the Pfizer vaccine in the 5-11 age group on: 19 October, 2 November, 9 November, 23 November, 30 November, 7 December, and 14 December.

- a. **The direct and indirect impacts on children.** Children who have COVID-19 will commonly have few or only mild respiratory symptoms. COVID-19 in this age group is rarely severe or fatal [18, 19], and the rate of severe COVID-19 disease in this age group is the lowest of any age group. However, there is a very small but real risk of MIS-C (described above) at this age which has occurred more frequently among ethnic minorities in the US [12, 20]. A very small proportion of children also experience persistent illness and ongoing symptoms, though evidence about its incidence is limited.
- b. In the current Delta outbreak in New Zealand (data to 19 November 2021), children aged 5-11 made up 14.9% of cases (1,003/6,714). Eight of these children were hospitalised but none were admitted to ICU. Of those who were hospitalised, all but one had a pre-existing condition and three were in hospital for less than six hours. As a comparison, between 16 June and 13 November 2021 in Sydney, 14,154 cases (19.4%) were aged 0-11 years and not eligible for vaccination. Of these cases, 632 were hospitalised, 9 were in ICU, and 0 patients died. It was not mentioned whether any of these cases had pre-existing conditions or comorbidities. The Sydney data further demonstrates that COVID-19 is relatively mild in most young children as despite accounting for 19.4% of cases in Sydney since 16 June, they account for only 5.9% of hospitalisations, 0.6% of ICU admissions, and no deaths [40].
- c. Children living with pre-existing conditions or comorbidities, from disadvantaged backgrounds, or those living within a lower socioeconomic status have a greater risk of severe disease from COVID-19 [12-15].
- d. Even though the direct effects of infection are generally less severe in children, this should not diminish the significance for those who have experienced worse outcomes [19]. Alongside the direct risks and impacts to health and individuals, COVID-19 also has indirect impacts for children on mental health, wellbeing, education and social development, and these are worsened by lockdowns and school closures [12, 21-23].
- e. **Children do play a role in transmission however it is significantly smaller than for adults.** Transmission within education settings has occurred but is limited and is more likely to occur between adults [24-26]. Transmission in households is much more common [27, 28]. The benefit of vaccination on onward transmission in households could be lower than in other settings due to the ongoing and close nature of exposure [29, 30], but this is not confirmed. The effect of vaccination of children on household transmission is unknown.
- f. **There are a number of equity considerations which are important to consider:**
 - i. Māori and Pacific children have been disproportionately affected in the current outbreak. To 19 November 2021, Māori made up 52% of cases in 5- to 11-year-olds, and Pacific children have made up 30% of cases among 5- to 11-year-olds.
 - ii. Māori and Pacific adults are at greater risk of COVID-19 hospitalisation and severe disease. Māori and Pacific adults have respectively 2.5-fold and 3-fold higher odds of being hospitalised compared to non-Māori, and Māori are likely to spend 4.9 days longer in hospital [31, 32].

- iii. Māori and Pacific children are more likely to live in multigenerational families housed in overcrowded conditions, increasing the risk of transmission. The younger age structure of the Māori population also means that a larger proportion are currently unable to be vaccinated and remain susceptible to infection and transmission, with a risk of onwards transmission to whānau and communities [33, 34], though the risk of transmission from children is lower than from adults.
 - iv. The vaccine rollout in adults resulted in inequities for Māori and Pacific adults, and the rollout for Māori and Pacific children aged 5-11 will need close consideration and more tailored implementation. This emphasises the need for culturally appropriate messaging and Māori-led initiatives. Whānau-based approaches to the 5-11 rollout may also improve uptake among Māori adults.
 - v. According to a Horizon Research survey, 72% of those who care for 5- to 11-year-olds would allow their child to receive the COVID-19 vaccine, however this was lower among Māori caregivers at 51% [41]. However, we note that the Māori adult rate of uptake and the Māori childhood immunisation rates are much higher than 51%. Given this we believe with a correctly tailored programme, high rates of immunisation in tamariki Māori are achievable.
- g. **The paediatric formulation of the vaccine has been approved for emergency use and rolled out in the USA, Canada, and Israel.** The Advisory Committee on Immunization Practices (ACIP) made an interim recommendation for the emergency use of the Pfizer vaccine in children aged 5-11 years in the United States for prevention of COVID-19 [19, 35]. This was unanimously supported by the Committee. In making this recommendation, ACIP considered the importance of COVID-19 as a public health problem, as well as benefits and harms, parents' values and preferences, acceptability, feasibility, resource use, and equity for use of the vaccine among children [19, 35]. ACIP additionally stated: "children from racial and ethnic minority groups have experienced a disproportionately high incidence of COVID-19 as well as secondary impacts of the COVID-19 pandemic such as reduced in-person learning"[19]. These comments have high relevance for New Zealand given the similar effects of the pandemic on Māori and Pacific peoples as described above.
- h. **In Australia, the Therapeutic Goods Administration (TGA)** provisionally approved the Pfizer vaccine as safe and effective for use among this age group on 5 December [36]. ATAGI recommends all 5–11-year-olds be vaccinated with an 8-week interval between doses, and that those at risk of severe disease, Aboriginal and Torres Strait Islanders, and children in crowded conditions or outbreak areas be prioritised [37].
- i. **Data are still accumulating from the real-world rollout of vaccines in 5- to 11-year-olds, and there is currently limited safety data available post-second dose.** Some adverse events in other age groups (e.g. myocarditis) have only become apparent following widespread rollout, and as noted above the trials in young children are too small to be able to detect rare side effects. Further data on potential side effects from the vaccine rollout in this age group in other countries will become progressively available.

- j. **On coadministration and other vaccines**, there is limited evidence on the safety and immunogenicity of coadministration of the Pfizer vaccine with other vaccines in all populations, however based on first principles of vaccinology it is likely to be safe and effective, particularly in younger age groups.
- k. **The wider National Immunisation Schedule** has been facing challenges for some time with declining vaccination rates since before COVID-19, and are particularly marked for Māori and Pacific infants and children. Catch-up campaigns for the MMR, HPV and Tdap vaccines were further delayed by COVID-19 and lockdowns. There is a risk that rolling out the Pfizer vaccine in this age group could further adversely impact the wider immunisation programme through diverting public health resources. This could increase the risk of outbreaks of other infectious diseases. The risk of a significant measles outbreak is of particular concern once the international borders re-open. Vaccination rates are lowest among Māori, and therefore there are equity concerns that there will be greater risk in this population. However, there is also the opportunity to increase coverage with other vaccines with a thoughtfully implemented COVID-19 vaccination programme in this age group.
- l. **On dosing intervals**, there are no data available about extending the interval between doses of the paediatric formulation of the Pfizer vaccine, however, emerging data in adults suggests that the immune response is likely improved by extending the dosing interval [42, 43]. This is consistent with basic principles of vaccinology and immunology which suggests that immune responses are generally better with longer intervals. There may also be a connection between shorter intervals and increased reactogenicity or adverse events, and one pre-print paper on individuals aged 12 and over has shown a statistically significant increase in myocarditis if the second dose was given at a shorter interval of less than 30 days [44]. Australia and Canada have recommended an 8-week interval between doses for 5-11-year-olds, noting this may improve immunogenicity and reduce side effects. Having a longer interval would also allow greater time to monitor international safety data.
- m. **On vaccine requirements**, there is a significant risk that use of vaccination mandates or certificates in this age group will result in exclusion and an inability to fully participate in schooling and extracurricular activities. This is likely to inequitably impact communities who are already experiencing disadvantage and where current vaccine coverage is poor. Concerns regarding possible stigmatisation and exclusions could be addressed in ways that do not necessarily influence the decision to use. For example, there could be a policy decision that children cannot be denied access to locations/events on the basis of vaccination status, which could be operationalised by not issuing vaccine certificates for this age group.

19. **CV TAG recommended:**

- a. **Two doses of the paediatric Pfizer vaccine be offered to all 5-11-year-olds in Aotearoa New Zealand, with an 8-week interval between doses.**

- b. Māori and Pacific children, children with high-risk pre-existing conditions, and children living with vulnerable people should be prioritised for vaccination and tailored programmes developed.
- c. On the schedule between doses:
 - i. The interval between doses can be shortened in limited circumstances to a minimum of 3 weeks, such as prior to the initiation of significant immunosuppression or international travel.
 - ii. Children who turn 12 after their first dose should follow the authorised schedule which uses the paediatric primary formulation (10 µg). They should not be offered the adolescent/adult formulation (30 µg) of the Pfizer COVID-19 vaccine.
 - iii. Children in this age group who experience a clinically significant adverse event after their first dose should be carefully reviewed by a specialist clinician. An individual risk:benefit assessment should be made on whether to administer the second dose. Children in this age group are not obliged to receive a second dose if not clinically appropriate.
- d. The paediatric Pfizer vaccine can be administered before, after, or at the same time as other vaccines in this age group.
- e. The adolescent/adult Pfizer vaccine formulation (30 µg) should not be used in children aged 5-11 years.
- f. Mandates, vaccine certificates or vaccine targets **must not** be used or required for this age group, and children in this age group should not be denied access to locations or events based on their vaccination status. There should be no unintended consequences in terms of participation if children in this age group are not vaccinated, and any use of mandates, certificates or targets that may formally or informally encourage inappropriate exclusion from activities. Exemptions from vaccination should therefore also not be required for this age group. We recommend specific public education campaigns about why children should not be excluded from activities, in order to reduce the risk of informal exclusions.
- g. Specific consideration must be given to promoting and improving vaccine access to groups that have experienced disproportionate COVID-19 morbidity and mortality, as well as those with barriers to routine health care, especially for Māori and Pacific peoples. This could be achieved through using the broad geographic accessibility of pharmacies and expanding school-focused strategies. Whānau centred approaches should be considered within these environments to improve primary vaccination and booster rates in the adult population.
- h. Emphasis must be given to using the rollout of the COVID-19 vaccine as an opportunity to improve delivery and uptake of the wider National Immunisation Schedule, and large-scale events with whānau-based approaches should be organised to aid catch-up campaigns for other vaccines. The coverage of the childhood National Immunisation Schedule should be closely monitored to ensure that the COVID-19 vaccination rollout for this age group does not adversely impact on the uptake of other important childhood vaccines.

- i. In making vaccination available, it should not be solely relied upon and other public health measures in schools and other educational settings should be strengthened, including ensuring good ventilation and filtration of air indoors, use of masks, physical distancing, and promotion of children staying at home if sick.


20. CV TAG will continue to monitor all relevant information (including vaccine efficacy data against emerging variants of concern and emerging evidence on the duration of immunity) and will update their recommendations as further evidence becomes available.

- a. New Zealand and international safety data will be carefully monitored, and the recommendations here will be reassessed by CV TAG in February 2022 prior to second doses being given to any 5–11-year-olds in Aotearoa New Zealand.
- b. Advice for severely immunocompromised children who may need a third primary dose will be reconsidered once further evidence emerges on the need, safety, and efficacy.

Recommendations

It is recommended that you:

2.	Note this advice has been received.	Yes/No
----	-------------------------------------	--------



Dr Ian Town
**Chief Science Advisor and
 Chair of the COVID-19 Vaccine Technical Advisory Group**

Signature _____

Dr Ashley Bloomfield

Director-General of Health

Date: